

Decoherent histories quantum mechanics with one real fine-grained history

Murray Gell-Mann^{1,*} and James B. Hartle^{1,2,†}

¹*Santa Fe Institute, Santa Fe, New Mexico 87501, USA*

²*Department of Physics, University of California, Santa Barbara, California 93106-9530, USA*

(Received 5 December 2011; published 25 June 2012)

Decoherent histories quantum theory is reformulated with the assumption that there is one “real” fine-grained history, specified in a preferred complete set of sum-over-histories variables. This real history is described by embedding it in an ensemble of comparable imagined fine-grained histories, not unlike the familiar ensemble of statistical mechanics. These histories are assigned extended probabilities, which can sometimes be negative or greater than one. As we will show, this construction implies that the real history is not completely accessible to experimental or other observational discovery. However, sufficiently and appropriately coarse-grained sets of alternative histories have standard probabilities providing information about the real fine-grained history that can be compared with observation. We recover the probabilities of decoherent histories quantum mechanics for sets of histories that are recorded and therefore decohere. Quantum mechanics can be viewed as a classical stochastic theory of histories with extended probabilities and a well-defined notion of reality common to all decoherent sets of alternative coarse-grained histories.

DOI: [10.1103/PhysRevA.85.062120](https://doi.org/10.1103/PhysRevA.85.062120)

PACS number(s): 03.65.Ta, 03.65.Yz, 98.80.Qc, 98.80.Bp

I. INTRODUCTION

Decoherent histories quantum mechanics¹ (DHQM) is logically consistent, in agreement with experiment as far as is known, applicable to cosmology, consistent with the rest of modern physics including special relativity and quantum field theory, and generalizable to include quantum gravity.² It is a framework for quantum cosmology and for understanding large scale features of the quantum universe ranging from the approximate applicability of classical physics under suitable conditions to the number of e-foldings of inflation. It includes the Copenhagen quantum theory of laboratory experiment as an approximation adequate for measurement situations. It is the only presently available formulation of quantum theory with all these properties.

DHQM assigns probabilities to the members of decoherent sets of alternative coarse-grained histories of the universe. By decoherent we mean that, as a consequence of the quantum state and dynamics of the universe, there is negligible quantum interference between coarse-grained histories in the set. A coarse-grained history can be regarded as a class of fine-grained ones. Fine-grained histories describe the system as completely as possible. Feynman paths for a particle are an example. Decoherent sets of alternative coarse-grained histories are called realms.

As usually formulated, DHQM presents two obstacles to the idea that there is one unique, real, fine-grained history of the universe that we experience at a highly coarse-grained level. The first is that, except for trivial cases, there are no decoherent sets of completely fine-grained histories. Negligible interference requires coarse graining such that the phases between

histories are washed out. The second obstacle is that there are realms that are mutually incompatible in the sense that there is no common finer-grained realm of which they are both coarse grainings. There appears to be no connection between the real history of one realm and that of an incompatible one.

In this paper we show how to overcome these obstacles. We overcome the first one by extending the notion of probability to include values outside the range $[0, 1]$. Extended probabilities can then be assigned to any set of alternative histories, in particular to fine-grained ones. We overcome the second obstacle by restricting the allowed sets of histories to those describable in a preferred set of variables—those used in a sum-over-histories treatment of quantum theory. The result will be a formulation of quantum theory that at first sight appears to be both an extension and a contraction of DHQM. It is an extension in that it uses extended probabilities; it is a contraction in the sense of using only a preferred set of variables. However, we shall see that in realistic situations it is equivalent to a sum-over-histories formulation of DHQM.

There are two basic starting points for this formulation. The first is the ensemble method of Gibbs [6], which has proved to be essential for describing the coarse-grained regularities of physical systems about which we have little fine-grained information, as in statistical mechanics. The same method is also useful for characterizing the complexity of such regularities [7]. The second starting point is the notion of extended probability [8].

The statistical mechanics of a classical gas of N particles in a box illustrates the ensemble method. The gas is described at a moment of time t_0 by giving the positions and momenta of the N particles; that is, by giving a point, $z_0 = z(t_0)$, in the $6N$ -dimensional phase space of the system. This point in phase space evolves in time; $z(t)$ follows Newton’s deterministic laws. That is the real fine-grained history of the gas. The evolution by a deterministic law is a regularity of the fine-grained history, but not one that is completely accessible or useful to us when N is large. There is no practical hope of measuring, storing, retrieving, or computing with all the information

*mgm@santafe.edu

†hartle@physics.ucsb.edu

¹For broad accounts of decoherent histories and related formulations, with references to important earlier literature, see, for example, [1–3]. For a short tutorial see [4].

²For the extension to quantum spacetime see, for example, [5].

involved in describing these regularities. The accessible, useful regularities, such as those summarized by the Navier-Stokes equation, are very coarse-grained. To describe them we conceptually embed the one real history of the particles in an ensemble (a set with probabilities) of imagined comparable fine-grained histories with various different initial conditions. The term ensemble indicates that probabilities are assigned to these initial conditions in such a way that the coarse-grained regularities have high probability. For example, one might consider a time-dependent coarse graining based on hydrodynamic quantities with probabilities representing local equilibrium. The phenomenological equations of classical hydrodynamics then hold with high probability.³ The assumption that the one real history is typical in this ensemble allows us to use the probabilities to bet on what will really happen in the future and what really happened in the past.

We will show how the quantum mechanics of a closed system can be formulated in a similar way. We assume the preferred variables of sum-over-histories quantum mechanics are the ones to be used for describing fine-grained histories.⁴ The one real fine-grained history of the universe is embedded conceptually in an ensemble of alternative fine-grained histories, each of which is assigned an *extended* probability [8] based on an assumed quantum state and Hamiltonian for the system. Extended probabilities obey the usual rules of probability theory except that they can be negative or greater than one for an alternative for which it cannot be determined whether it occurs or not (as on the alternative histories in the two-slit experiment). From the extended probabilities of fine-grained histories, extended probabilities can be constructed for sets of coarse-grained alternative histories that are classes of fine-grained histories. Sufficiently and appropriately coarse-grained ensembles have only probabilities between 0 and 1 and can therefore be used to bet on the outcomes of experiments that test the theory. The result, as mentioned in Ref. [8], is that quantum mechanics can be viewed as a classical stochastic theory with extended probabilities.

The Bell inequalities together with the experiments verifying their violation restrict the nature of any classical stochastic theory that might reproduce the probabilities of quantum mechanics. Either locality or positive probabilities or both must be given up (see, e.g., [13]). As we will show quantum mechanics naturally allows extended probabilities which can be nonpositive. It is with these that we construct the ensemble of fine-grained histories. Extended probabilities are therefore an essential part of our story.⁵

This formulation of the quantum mechanics of closed systems is not a replacement for decoherent histories quantum

theory but rather a different starting point for it. As we will see, we recover a sum-over-histories version of DHQM in the end for realistic situations.

Our paper is organized as follows: Section II explains how quantum mechanics can be formulated as a prescription for the extended probabilities of an ensemble of fine-grained histories containing one real history. Section III discusses the role of records and the connection to decoherent histories quantum theory. Section IV discusses the implications of this formulation of quantum mechanics. Section VI contains a summary and further discussion of the implications.

II. THE QUANTUM ENSEMBLE OF FINE-GRAINED HISTORIES

This section formulates the quantum mechanics of a closed system as a theory of one real fine-grained history embedded in an ensemble of comparable alternative fine-grained histories described in a preferred set of variables and assigned extended probabilities. To keep the discussion manageable, we neglect gross fluctuations in the geometry of spacetime. We then in effect consider a system of a box of particles or fields moving in a fixed, presumably expanding, background spacetime. Well-defined notions of space and time are then available, as well as the usual machinery of quantum theory—amplitudes, operators, a Hilbert space of states, etc. The important thing is that the system is closed so that observers and measuring apparatus (if any) and all other physical systems are within the box and are part of the quantum system being described. We can think of this as a simplified model of the universe.

The fundamental theory of this closed system consists of two parts: the system's Hamiltonian H specifying the dynamics (assumed independent of time for simplicity) and the quantum state specified by a wave function Ψ .

A. Four ingredients of the formulation

This formulation is specified by four ingredients:

- (1) The preferred set of variables in terms of which the one real fine-grained history is described, as well as the alternative fine-grained histories of the ensemble in which the one history is embedded.
- (2) An extended notion of probability that reflects the notion of ignorance in quantum mechanics.
- (3) The prescription that assigns extended probabilities to the members of the ensemble of alternative fine-grained histories using the system's quantum state and Hamiltonian.
- (4) Coarse graining of the fine-grained set of histories leading to coarse-grained sets that can be recorded and decoherent and can have standard probabilities.

We now discuss these in turn:

Preferred variables. The fine-grained histories are described by a preferred set of variables which we take to be those of a sum-over-histories formulation of quantum mechanics. They are histories of particle positions in the case of particles, four-dimensional field configurations—both bosonic and fermionic—in the case of quantum field theory, and histories of geometries and fields in the case of semiclassical quantum gravity. Histories of these variables are assumed to be the most refined description of the system possible.

³See, for example, [9] for an exposition in the present context.

⁴This assumption fits with the idea that the sum-over-histories formulation of quantum theory may be a more general and therefore a more fundamental framework for quantum mechanics [10,11]. Extensions of usual quantum theory to incorporate quantum spacetime seem naturally formulated in this way (see, e.g., [5,12]).

⁵Bohm theory [14] is an example of a version of quantum mechanics that assigns usual probabilities to fine-grained histories of particle positions but gives up on locality in the sense it is used in the Bell inequality discussion.

A strong case can be made that these histories are adequate for the prediction of all observable quantities [15]. Particle momentum, for instance, can be defined in terms of histories of position in a time-of-flight setup that would measure velocity. We do not rule out introducing operators, transformation theory, etc. at some later stage but we begin by assuming this preferred set of variables for fine-grained histories.

To describe these preferred variables we assume a particular Lorentz frame and let t be the time coordinate of that frame. We denote the preferred variables by q^i or just q for short. For particles, i might be x, y, z and a particle label. For fields, i would include the label \vec{x} of the spatial point. We denote the configuration space spanned by q^i by \mathcal{C} . A fine-grained history is a path $q(t)$ in \mathcal{C} that we assume to be single-valued—one and only one value of q for each t . The set of all fine-grained histories between an arbitrary pair of times t_0 and t_f is the set of all such paths $\{q^i(t)\}$ between these two times. They are continuous but typically nondifferentiable.

Extended probabilities. Probabilities can be usefully understood as instructions for making fair bets [16]. To hold that the probability of an alternative α is $p(\alpha)$ means the following: Suppose that there is a bet on whether α occurs with payoff S_α (of either sign) if it does. You will put up $p(\alpha)S_\alpha$ and consider it a fair bet. Probabilities can thus be said to express our ignorance with respect to whether the alternative α occurs.

All the usual rules of probability theory, including the restriction of values to the range $[0,1]$, follow from the requirement that a bookie not be able to offer you a “Dutch book” in which you will put up the stake $p(\alpha)S_\alpha$ but be guaranteed to always lose, not just on average, but each time the bet is made [16,17].

Implicit in the above definition of probability in terms of fair bets is the assumption that it can be settled whether the alternative α occurs or not. Elementary physics assumes that any alternative that can be described can be determined without significantly affecting its value. Every alternative in elementary physics is therefore in principle the basis of a settleable bet, however difficult it may be to settle it in practice. For instance, we assume that the value of the initial condition z_0 of the box of gas discussed in the preceding section is in principle the basis for a settleable bet even though it is impossible in practice to determine z_0 . Thus, when we construct an ensemble reflecting our ignorance of z_0 , we assign probabilities obeying the usual rules to the different values it might take.

But in usual quantum theory there are alternatives that can be described but are not the basis for settleable bets. A classic example is provided by the two-slit experiment illustrated in Fig. 1. A bet on whether the electron went through the upper slit or the lower slit is not settleable without carrying out a measurement that would significantly disturb the system. Alternatives that are not the basis of settleable bets are a new kind of ignorance not found in elementary physics and they can be usefully described by extending the classical notion of probability. In Ref. [8] we proposed the following simple extension: Keep the rules of probability as they are, but allow the values to be outside the range $[0,1]$ for alternatives corresponding to nonsettleable bets. Evidently, this extension does not lead to Dutch books because no book of any kind can be made on nonsettleable alternatives.




FIG. 1. The two-slit experiment. An electron gun at left emits an electron traveling toward a screen with two slits; its progress in space recapitulating its evolution in time. The electron is detected at a further screen at a position y with a probability density that exhibits an interference pattern. A coarse-grained set of histories for the electron is defined by specifying the slit (U or L) through which the electron passes through and ranges Δ of the position y where it is detected. In the absence of the record of a measurement it is not possible to settle a bet on the which of these histories occurred.

Even one out-of-range probability in an exhaustive set of exclusive alternatives means that there are no settleable bets on which the alternative occurs. If there were, a Dutch book could be constructed using only the alternatives that have out-of-range extended probabilities. The excursion out of the normal range $[0,1]$ is a measure of how much coarse graining is needed to find alternatives that have probabilities in the normal range [8,18].

The quantum history ensemble will consist of alternative fine-grained histories assigned extended probabilities. We now turn to how the ensemble is constructed.

The fundamental distribution. To complete the formulation of quantum theory starting with an ensemble of fine-grained histories, we need to specify the fundamental distribution $w[q(t)]$ that assigns an extended probability⁶ to each history $q(t)$. This formula will depend on the initial quantum state of the box represented by a Schrödinger picture wave function $\hat{\Psi}(q, t_0)$. (Hats denote the Schrödinger picture.) This is just as much a part of the necessary theoretical structure for prediction in the universe as the Hamiltonian⁷ H . There are no quantum-mechanical predictions of any kind that do not depend on both. The Schrödinger picture wave function evolves in time according to

$$\hat{\Psi}(q, t) = e^{-iH(t-t_0)/\hbar} \hat{\Psi}(q_0, t_0). \quad (2.1)$$

To keep the discussion manageable, assume that all coarse-grained alternatives lie between an initial t_0 and a final time t_f . That is not a loss of generality since t_f can be as large as desired. Elementary causality shows that we need not consider fine-grained histories at times later than t_f (see, e.g., [20],

⁶The distribution is a probability functional density for continuous q , but we rely on the reader to make this qualification where appropriate.

⁷In more general quantum gravitational contexts a theory of the quantum state like the one that yields the no-boundary wave function [19] may derive the state from the dynamical theory.

Sec. VII C). Thus, we consider single-valued histories $q(t)$ that take initial values q_0 at $t = t_0$ and final values q_f at $t = t_f$. We then postulate the fundamental distribution of extended probabilities is [8]

$$w[q(t)] \equiv \text{Re}[\hat{\Psi}^*(q_f, t_f) \exp\{i\mathcal{S}[q(t)]/\hbar\} \hat{\Psi}(q_0, t_0)], \quad (2.2)$$

where $\mathcal{S}[q(t)]$ is the action functional corresponding to the Hamiltonian H . We will show in Sec. III that this formula reproduces the probabilities of DHQM when there are records of histories that can be used to settle bets on which history actually occurs. At this point in the exposition, however, (2.2) is a postulate.

So defined, the extended probabilities sum to one since the integral over all paths from t_0 to t_f (including over q_0 and q_f) is

$$\int \delta q w[q(t)] = \int dq_f \hat{\Psi}^*(q_f, t_f) \hat{\Psi}(q_f, t_f) = 1, \quad (2.3)$$

because the sum over all paths just evolves the wave function at t_0 to t_f to give (2.3).

The distribution $w[q(t)]$ defined by Eq. (2.2) will have values outside the range $[0,1]$ for some $q(t)$. To see that, suppose there is a history $q(t)$ with $w[q(t)]$ within the range $[0,1]$. Then a variation of $q(t)$ that leaves q_0 and q_f unchanged can still contribute significantly to the action in Eq. (2.2) and change the sign of $w[q(t)]$ from positive to negative. As we will see in Sec. III, these negative extended probabilities mean that the set of fine-grained histories is not the basis for a settleable bet on what the real fine-grained history is like.

Coarse graining. The set of alternative fine-grained histories $\{q(t)\}$ can be coarse grained by partitioning it into an exhaustive set of exclusive classes c_α with $\alpha = 1, 2, \dots$. The extended probability of the class c_α is the sum of the extended probabilities of all its members; that is,

$$p(\alpha) = \int_{c_\alpha} \delta q w[q(t)], \quad (2.4)$$

where the sum is over all $q(t)$ in the class c_α , including a sum over q_0 and q_f .

Sufficient coarse graining will lead to sets with all positive probabilities if only because the completely coarse-grained set consisting of all histories has probability one.⁸ Suitably, coarse grained sets can be recorded and decoherent and can have standard probabilities. They therefore describe alternatives that are the basis of settleable bets. We discuss this in more detail in the next section, but we conclude this one by introducing useful operator representations of most of our formulas.

B. Operators

Using (2.2), the expression (2.4) for the extended probability of a coarse-grained history c_α can be written more explicitly as

$$p(\alpha) = \int dq_f \int dq_0 \int_{[q_0 c_\alpha q_f]} \delta q \times \text{Re}[\hat{\Psi}^*(q_f, t_f) \exp\{i\mathcal{S}[q(t)]/\hbar\} \hat{\Psi}(q_0, t_0)], \quad (2.5)$$

⁸See [8] for a more quantitative discussion of how coarse graining leads to usual probabilities.

where the notation $[q_0 c_\alpha q_f]$ means that the sum is over all histories in the class c_α that begin at q_0 and end at q_f . This can be conveniently written in operator form as

$$p(\alpha) = \text{Re}[\langle \hat{\Psi}(t_f) | \hat{C}_\alpha | \hat{\Psi}(t_0) \rangle], \quad (2.6)$$

where the matrix elements of the class operator \hat{C}_α in the Schrödinger picture are defined by

$$\langle q_f | \hat{C}_\alpha | q_0 \rangle \equiv \int_{[q_0, c_\alpha, q_f]} \delta q e^{i\mathcal{S}[q(t)]/\hbar}, \quad (2.7)$$

and $\hat{\Psi}(q, t) = \langle q | \hat{\Psi}(t) \rangle$.

The formulas simplify even further in the Heisenberg picture. Referred to the time t_0 , the Heisenberg picture state is

$$|\Psi\rangle \equiv e^{iH(t_f - t_0)/\hbar} |\hat{\Psi}(t_0)\rangle. \quad (2.8)$$

The equivalence between sum-over-histories evolution and Hamiltonian evolution shows that the Heisenberg picture state $|\Psi\rangle$ is constant in time. Then, if we define

$$C_\alpha \equiv e^{iH(t_f - t_0)/\hbar} \hat{C}_\alpha, \quad (2.9)$$

we find⁹

$$p(\alpha) = \text{Re}[\langle \Psi | C_\alpha | \Psi \rangle]. \quad (2.10)$$

Extended probabilities satisfy all the rules of probability theory except the requirement that probabilities lie in the range $[0,1]$. In particular, they must satisfy the usual sum rules connecting the extended probabilities for coarse-grained sets related by operations of fine and coarse graining. Suppose that $\{\bar{c}_\alpha\}$ is a coarse graining of a set $\{c_\alpha\}$. This means that the even coarser-grained set is a partition of the finer-grained set into an exhaustive set of exclusive classes so that

$$\bar{c}_\alpha = \bigcup_{\alpha \in \bar{\alpha}} c_\alpha. \quad (2.11)$$

The rules of extended probabilities require the sum rules

$$\bar{p}(\bar{\alpha}) = \sum_{\alpha \in \bar{\alpha}} p(\alpha). \quad (2.12)$$

These are satisfied exactly as a consequence of (2.4), which leads to

$$\bar{C}_\alpha = \sum_{\alpha \in \bar{\alpha}} C_\alpha. \quad (2.13)$$

A familiar example of a coarse graining of a fine-grained set of histories concerns the motion of a particle in one dimension x . The set of fine-grained paths can be partitioned into classes by using exhaustive sets of exclusive intervals of x , $\{\Delta_{\alpha_k}^k\}$, with $\alpha_k = 1, 2, \dots$, at a sequence of times t_k $k = 1, 2, \dots, n$. The index k allows for the sets to be different at different times. The index α_k labels the interval in each set. The alternatives at each time correspond to an exhaustive set of exclusive (Schrödinger

⁹This is the expression used by Goldstein and Page [21] to define their linear positivity condition that restricts sets of coarse-grained histories to those for which (2.10) is positive. The fundamental distribution (2.2) is a related expression that assigns extended probabilities to fine-grained histories without restriction.

picture) projection operators $\{\hat{P}_{\alpha_k}^k\}$ onto the intervals $\Delta_{\alpha_k}^k$. A coarse-grained history α is defined by a particular sequence of intervals at the possible times, $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_n)$. The class c_α consists of the bundle of paths $q(t)$ that pass through these intervals at the assigned times. The class operator in the Heisenberg picture defined by Eq. (2.7) can be shown (see, e.g., [22]) to be given by

$$C_\alpha = P_{\alpha_n}^n(t_n) \cdots P_{\alpha_1}^1(t_1). \quad (2.14)$$

The extended probabilities of these histories are given by (2.10); some explicit calculations can be found in Ref. [18].

III. SETTLEABLE BETS, RECORDS, AND DECOHERENCE

A. Utilization by information gathering and utilizing systems

In this section we turn to the question of how extended probability ensemble quantum mechanics (EPE) is used. Information gathering and utilizing systems (IGUSs) like ourselves exploit the regularities summarized by physical theories to construct schemata, make predictions, and direct behavior [3]. Even a frog aiming to catch a fly can be said to be exploiting a rudimentary classical approximation to quantum mechanics (perhaps hard wired). The frog is in effect betting on these regularities.

We can imagine an IGUS equipped with the EPE formulation of quantum theory of the previous section and theories of the quantum state $\hat{\Psi}$ and action \mathcal{S} . Such an IGUS might use the theory plus data acquired in previous observations to make bets that will be settled by future data it may acquire. That is true whether or not the bet concerns the outcome of a delicate laboratory experiment, the outcome of the next election, what happened in the distant past, or whether the universe will be destroyed in a big crunch singularity in the far future.

To make a bet the IGUS must first specify the alternatives that are the basis for the bet, and then prescribe the means to settle it. The only candidates for settleable bets are sets of alternative histories that are sufficiently coarse-grained that all the $p(\alpha)$ are positive so that the set has genuine probabilities. Of course, even with that restriction there is an infinity of possibilities.

B. Records and decoherence

Among the wealth of possibilities for alternative histories to bet on and for ways of settling such bets, one general property stands out: Bets are settled using records of the alternatives one is betting on. A set of histories $\{C_\alpha\}$ is recorded if there are alternatives with projection operators $\{R_\alpha\}$ at one time that are correlated with the histories. These alternatives need not be ones that are used or even could be used by IGUSs. As in the examples of Joos and Zeh [23], there is a record of the position of a dust grain in interstellar space in all the CMB photons that have scattered from it and left the vicinity at the speed of light.

A very general notion of record is the following: We say that a set of histories $\{C_\alpha\}$ is (strongly) recorded if there is an exhaustive set of orthogonal exclusive projection operators $\{R_\alpha\}$ on regions of the configuration space spanned by the q^i

at a time after the last alternative in C_α such that

$$R_\alpha|\Psi\rangle \approx C_\alpha|\Psi\rangle \quad (3.1)$$

to a suitable approximation appropriate for realistic situations.¹⁰ The projection R_α defines the record of the history C_α . Some approximation must be allowed in Eq. (3.1) because we cannot expect realistic records to be *exactly* correlated with the histories they record. The degree of approximation is treated in Eq. [8]. We now discuss how some familiar features of quantum mechanics are recovered for sets of histories $\{C_\alpha\}$ that are recorded in the sense of Eq. (3.1).

From the fundamental distribution (2.10) and the definition of recorded sets of histories (3.1) it follows that

$$p(\alpha) \approx \langle \Psi | R_\alpha | \Psi \rangle = \langle \Psi | R_\alpha C_\alpha | \Psi \rangle \approx \|C_\alpha|\Psi\rangle\|^2. \quad (3.2)$$

The extended probabilities are therefore positive for a set of histories for which there are records. They are genuine probabilities with which to make fair bets, and the records supply a way to settle each bet. Further, the values of the probabilities are given by the usual quantum-mechanical square of an amplitude (the branch state vector).

A recorded set of histories is decoherent. That is because the projections R_α are orthogonal, so that

$$0 = \langle \Psi | R_\alpha^\dagger R_\beta | \Psi \rangle \approx \langle \Psi | C_\alpha^\dagger C_\beta | \Psi \rangle \approx 0, \quad \alpha \neq \beta, \quad (3.3)$$

which is the condition for (medium) decoherence.¹¹ The sets of histories that are the basis for bets settleable by records are decoherent and have standard probabilities.

C. EPE as a version of DHQM

In equations (3.3) and (3.2) we have the central relations of decoherent histories quantum mechanics. Equation (3.3) is the condition for decoherence of the set of coarse-grained histories represented by the $\{C_\alpha\}$, and (3.2) gives their standard probabilities. Thus we have recovered a sum-over-histories version of decoherent histories. The probabilities of EPE that satisfy the sum rules of probability theory exactly are recovered only to the extent that the histories are recorded as in Eq. (3.1). The DHQM probabilities will satisfy these rules approximately, but in realistic situations there is no physically significant difference between the two sorts of probability. Extended probability ensemble quantum mechanics is a version of decoherent histories quantum mechanics.

Expositions of EPE and traditional DHQM (TDHQM) begin at the same starting point and end with essentially the same predictions for probabilities of histories in realms. But the routes to these ends differ—in particular with regard to the role played by records and decoherence. It may be helpful to the reader to compare these different routes.

¹⁰There are other general notions of record. In the terminology used in Ref. [8] we are defining strong records. There a further discussion of the appropriate degree of approximation can be found.

¹¹In a general operator context the decoherence condition implies that records of histories exist, but not necessarily in a preferred set of variables as we are assuming here. For recovering DHQM we need only that records imply decoherence, not the other way around.

Both approaches to DHQM aim at the probabilities for sets of alternative histories represented by $\{C_\alpha\}$. Both start from the observation that there are interfering sets of histories that cannot consistently be assigned probabilities.

TDHQM assigns probabilities only to sets of histories for which quantum interference between the members of the set is negligible as a consequence of the state $|\Psi\rangle$ and Hamiltonian H . TDHQM does not treat sets of histories for which interference is non-negligible.

Decoherence means that the branch state vectors $\{|\Psi_\alpha\rangle\}$ are nearly mutually orthogonal. This means that records of the histories exist if only trivially as projections $\{R_\alpha\}$ on the branch state vectors. The approximate probabilities of the histories are given by $p(\alpha) = ||C_\alpha|\Psi\rangle||^2$.

EPE starts from the idea that probabilities are instructions for settleable bets. But interfering sets of alternative histories are not the basis of a settleable bet. The notion of probability is extended to give extended probabilities to nonsettleable alternatives.

Records of histories are defined as the means of settling bets on which of a set of alternative histories occurs. Decoherence is deduced as a consequence of a strong notion of records. In contrast to records following from decoherence in TDHQM, in EPE decoherence follows from records. The probabilities of recorded histories in EPE are $p(\alpha) = \text{Re}\langle\Psi|C_\alpha|\Psi\rangle$, which differ negligibly from the TDHQM $p(\alpha) = ||C_\alpha|\Psi\rangle||^2$ in realistic situations.

D. States and the quantum-mechanical arrow of time

The extended probability formulation of quantum theory elaborated in the previous section is time-neutral—incorporating no fundamental arrow of time. That can be seen from the fundamental distribution (2.10). Field theory is invariant under CPT transformations that reverse the time order of the alternatives in a C_α . [That is immediate in the case of chains of alternatives at moments of time, as in Eq. (2.14).] The result is to complex conjugate the matrix elements in Eq. (2.10) and (2.6), but the resulting extended probabilities $p(\alpha)$ are unchanged. At this level of generality there is no quantum-mechanical arrow of time as there is in usual quantum theory and no fundamental distinction between past and future.

That means that there is not a notion of a state at a moment of time that summarizes the past for prediction of the future. That is a time-asymmetric notion.¹²

However, the formula (3.2) for the probabilities of a recorded set of histories is not time-neutral. Written out for histories that are sequences of alternatives at definite moments of time [cf. (2.14)] it reads

$$p(\alpha) \approx ||P_{\alpha_n}^n(t_n) \cdots P_{\alpha_1}^1(t_1)|\Psi\rangle||^2. \quad (3.4)$$

The projections in the chain are time ordered.¹³ As already mentioned, that is not an essential asymmetry because the order

could be reversed by a CPT transformation. But, in any case, there is the state at one end of the chain and nothing at the other end. This asymmetry is what is called the quantum-mechanical arrow of time.¹⁴

How did we arrive at a formulation with a quantum-mechanical arrow of time from one that did not have one? The answer is that the notion of record introduced it. To see that, first note that a consequence of Eq. (3.1) is

$$R_\beta C_\alpha |\Psi\rangle \approx \delta_{\alpha\beta} C_\alpha |\Psi\rangle. \quad (3.5)$$

The combinations $R_\beta C_\alpha$ can be considered as a set of histories in which the time of the records t_{rec} is after the last time t_n in the chain in Eq. (3.4) ($t_{\text{rec}} \geq t_n$). The result is that the probabilities are

$$p(\beta, \alpha) \approx \delta_{\beta\alpha} p(\alpha), \quad (3.6)$$

which concisely expresses the correlation between records and history. With the usual conventions for past and future the R s then can be said to record the past.

The above discussion shows that when the records defined by Eq. (3.1) are considered as part of history they come at the opposite end of the chain from the state $|\Psi\rangle$. That is consistent with our intuitive understanding of the processes by which records are formed in a universe like ours that has an initial low-entropy state leading to a second law of thermodynamics. (See [9] for further discussion.)

The quantum-mechanical arrow of time of DHQM allows us to introduce a notion of state at a moment of time as we discussed in Ref. [24]. Suppose we have a decoherent set of histories consisting of sets of $(\alpha_n, \dots, \alpha_1)$ at sequence of times $t_1 < t_2 < \dots < t_n$ with $t_1 > t_0$. Suppose that at time t_k we know from present records that particular alternatives $(\alpha_k, \dots, \alpha_1)$ happened in the past. Predictions for future alternatives are given by the conditional probabilities

$$p(\alpha_n, \dots, \alpha_{k+1} | \alpha_k, \dots, \alpha_1) \equiv \frac{p(\alpha_n, \dots, \alpha_1)}{p(\alpha_k, \dots, \alpha_1)}, \quad (3.7)$$

where we have dropped the time labels to keep the notation concise. This can be rewritten

$$\begin{aligned} & p(\alpha_n, \dots, \alpha_{k+1} | \alpha_k, \dots, \alpha_1) \\ & \equiv ||P_{\alpha_n}^n(t_n) \cdots P_{\alpha_{k+1}}^{k+1}(t_{k+1})|\Psi_{\alpha_k \dots \alpha_1}\rangle||^2, \end{aligned} \quad (3.8)$$

where

$$|\Psi_{\alpha_k \dots \alpha_1}\rangle \equiv \frac{P_{\alpha_k}^k(t_k) \cdots P_{\alpha_1}^1(t_1)|\Psi\rangle}{||P_{\alpha_k}^k(t_k) \cdots P_{\alpha_1}^1(t_1)|\Psi\rangle||}. \quad (3.9)$$

The state $|\Psi_{\alpha_k \dots \alpha_1}\rangle$ defines a state at the time t_k that has evolved from the beginning by unitary evolution (constant

equations of motion) as there is in a general operator formulation (see, e.g., [25]).

¹⁴In a more general time-neutral formulation of decoherent histories quantum theory [26] this asymmetry can be seen as arising from an initial condition that is a pure state or a somewhat mixed state and a final condition of total indifference (a density matrix proportional to the unit matrix). Indeed from that perspective the present formulation can be seen as one way of justifying a final condition of total indifference.

¹²See, for example, the discussion in Ref. [24].

¹³In this paper expressions like (3.4) are to be understood as abbreviations for particular sum-over-histories expressions (in the preferred variables q^i) following from (2.2) and (2.4). Thus, there is no freedom to change the times $\{t_k\}$ and thereby the descriptions of the projections (to a different set of variables using the Heisenberg

in the Heisenberg picture) and projection (reduction). It summarizes past information for future prediction. We recover this time asymmetric notion of state at a moment of time, not generally, not exactly, but as a special feature of those sets of histories that are approximately recorded and are therefore approximately decoherent.¹⁵

IV. IMPLICATIONS OF THE EXTENDED PROBABILITY ENSEMBLE FORMULATION

In this section we develop the implications of the extended probability ensemble formulation of quantum theory (EPE) in which the “real” history of the universe is embedded in an ensemble of comparable imagined histories with extended probabilities.

Throughout we will assume that histories are specified in the variables of sum-over-histories quantum mechanics (e.g., quantum fields) whether they are real or imagined, fine grained or coarse grained. Notions like realms (decoherent sets of coarse-grained histories) are also to be understood in this sum-over-histories context.

A coarse graining is a partition of the histories of the ensemble into an exhaustive set of exclusive classes, which are the coarse-grained histories. The one real fine-grained history must lie in one of these classes. That class is the real coarse-grained history in the set. With sufficient and appropriate coarse graining we have realms with standard probabilities that can be the basis of settleable bets on what the real history in the set is like. One coarse-grained history in each realm is real.

By measurement and other observation we acquire data D on what the real coarse-grained history is in any realm in which D is valid in some histories but not in others. With these data we also acquire coarse-grained information on what the real fine-grained history is. As we make further observations we learn more and more about the real fine-grained history.

However, this process of progressive discovery of reality can never be carried to the completely fine-grained level. That is because the set of fine-grained histories is not decoherent, not recorded, and therefore not the basis for settleable bets.¹⁶

The ensemble formulation permits the straightforward and unqualified use of ordinary language in quantum-mechanical discussion. Statements about what happened, or is happening, or will happen refer to the real history in situations where it is possible to bet on records that check these. That is, we can straightforwardly use these words in discussing realms.¹⁷

Because of the restriction to sum-over-histories variables, all coarse-grained sets are coarse grainings of the unique

common fine-grained set of histories. However, even with this restriction there will generally be mutually incompatible realms (e.g., as discussed in Ref. [9]). That is, there are decoherent coarse-grained sets for which there is not a finer-grained decoherent set of which they are both coarse grainings.¹⁸

EPE is a four-dimensional presentation of quantum theory. The histories in the ensemble are spacetime histories. A spacelike surface of constant time in a particular Lorentz frame (of the kind we have assumed throughout) defines a notion of past, present, and future. For any realm, and any such spacelike surface, the unique real fine-grained history selects one coarse-grained history that gives a description of what is happening in the present, what did happen in the past, and what will happen in the future.

As is well known, spacetime formulations of theories are not inconsistent with notions of causality and IGUS-specific distinctions of past, present, and future (see, e.g., [29]). The physical structure of our universe makes it easier to acquire information about times toward the big bang (what we call the past) than about times away from the big bang (what we call the future). The fact that the sources of electromagnetic radiation that we receive now lie in the past is just one of the several reasons for this [29]. That is why there are more events in the past that can be retrodicted with high probability conditioned on present data than there are events in the future that can be predicted. Such time asymmetries are due to the particular quantum state of our universe.¹⁹

V. TYPICALITY RESTRICTIONS ON COARSE GRAININGS

Families of quasiclassical realms are striking features of our quantum universe. These are families of decoherent sets of coarse-grained alternative histories defined by quasiclassical variables—averages over suitable volumes of densities of approximately conserved quantities such as energy, momentum, and particle number. The histories consist largely of related but branch-dependent projections onto ranges of quasiclassical variables at a succession of times. Each history with a non-negligible probability constitutes a narrative, with individual histories exhibiting patterns of correlation implied by closed sets of effective equations of motion interrupted by frequent small fluctuations and occasional major branchings (as in measurement situations).

Human IGUSs make use of coarse grainings of a quasiclassical realm to describe the everyday world of experience—tables and chairs, stars and galaxies, measurements and their results. To be useful, a theory like that summarized in Eq. (2.2)

¹⁵Another property of ordinary quantum mechanics that emerges only for recorded history is the usual quantum-mechanical notion of nonentangled, noninteracting subsystems represented by product states. The extended probabilities generally do not factor into products, as noted by Diósi [27], but the probabilities for recorded histories do, if only approximately [8].

¹⁶The fine-grained histories of classical statistical mechanics are only in principle the basis of a settleable bet.

¹⁷This is in contrast with the usual formulation of DHQM where it is necessary to qualify the use of “happen” by choosing to which mutually incompatible family of realms it refers [28].

¹⁸Imagine for example a closed system of two kinds of particles: red and blue. It seems likely that a set of suitably coarse-grained histories following only the positions of the red particles could be decohered by the interactions with the blue particles. There could also be a set of coarse-grained histories following only the positions of the blue particles that are made to decohere by interactions with the red. These realms could be mutually incompatible, with no finer-grained decoherent set following the positions of both.

¹⁹For discussions in which the authors participated, see, for example, [9,26,29,30].

must reliably describe the coarse-grained regularities of the real history in a quasiclassical realm.

But what about other recorded coarse grainings that are allowed by the very general definition of record in (3.1)? It seems likely that there are recorded sets of coarse grained histories for which the ensemble defined by Eq. (2.2) will not give a reliable description.²⁰ The allowed coarse-grained sets must therefore be more restricted than those allowed by Eq. (3.1). We now describe how this works both in classical statistical mechanics and quantum mechanics.

A. Classical statistical mechanics

The ensembles of the classical nonequilibrium statistical mechanics of a box of gas are usually employed to make predictions of regularities defined by quasiclassical coarse grainings. (In the classical context we refer to these as hydrodynamic coarse grainings.) Hydrodynamic variables are readily accessible to observation. Indeed, the probabilities of these ensembles can be defined by maximizing the measure of ignorance (entropy) of fine-grained histories while holding fixed the ensemble averages of relevant hydrodynamic variables. Statistical mechanics based on these probabilities makes successful predictions for such hydrodynamic coarse grainings.

But there are coarse grainings for which the predictions of these ensembles could fail. Coarse grainings of classical histories can be defined by partitioning the space of the values of their initial phase space coordinates z_0 . Consider the class of coarse grainings in which this initial phase space is partitioned into a small region around some particular point z_1 and the rest of the space. The classical ensembles will typically assign a very low probability to the small region and a probability near unity to the rest.

Suppose the point z_1 happens to coincide with the initial condition z_r of the real history of the gas. Then the ensemble will have predicted a small probability for what really will happen and a probability near unity for something else happening.

Of course, the technology does not exist to explore such coarse grainings experimentally for a box with a large number of particles. Even if it did, it is unlikely that we would explore the one in which the real history has a small probability. Those limitations, however, should not obscure the fact that in statistical mechanics there is some coarse graining for which the real history has negligible probability. The probabilities assigned by the ensemble do not, in this sense, constitute a good description of the real history.

If such coarse grainings are not accessible by realistic experiment then the theory can perhaps be augmented by restrictions on the coarse-grainings for which it can be employed—to quasiclassical realms for instance. But if observations can be carried out to decide which history in such a coarse graining actually occurs then we have to allow that the observed (real) history may be one that was assigned a very low probability.

A characteristic feature of such coarse grainings is that the real coarse-grained history is not a typical history of the

ensemble. More precisely, suppose $\{p(\alpha)\}$ are the probabilities assigned by the ensemble to the alternative coarse-grained histories; let us call the real coarse-grained history r . Then the following typicality condition [7] is not satisfied:

$$-\log p(r) \ll -\sum_{\alpha} p(\alpha) \log p(\alpha) = S, \quad (5.1)$$

where S is the entropy of the ensemble's probability distribution.

Would we discard statistical mechanics if such an experiment were carried out with this result? It would be more sensible to restrict the coarse grainings that test the theory to ones in which the typicality condition (5.1) is satisfied. Such coarse grainings cannot be identified in advance of the determination of the real coarse-grained history. But typicality can be calculated when the real results are in and their significance as a test of the theory is assessed accordingly.

B. Quantum mechanics

The situation in the EPE formulation of quantum mechanics is similar. For any fine-grained history it is possible to find a coarse graining in which its probability is very small or even zero. For example, consider a partition of the fine-grained histories $q(t)$ by whether the values of q at a time t_1 are in a small region around q_1 or elsewhere. Unless the state and dynamics are very special, the fundamental distribution (2.2) will assign a small probability to the small region. If q_1 happened to coincide with the q of the real history at that time the real history would be assigned a small or zero probability.²¹

The notion of record in Eq. (3.1) that implies medium decoherence is general and mathematically simple. The quasiclassical coarse grainings used by human IGUSs are a much more restricted class. Restrictions on either the coarse grainings or their records can still lead to theories that are consistent with known observations and would rule out examples such as those discussed above.

Whatever the restriction on records and coarse grainings, one should check that the real coarse-grained history that emerges has a probability that is typical of the ensemble of coarse-grained histories before throwing out the theory in Eq. (2.2), which consists of $\hat{\Psi}$, S , and quantum mechanics itself.

VI. DISCUSSION

By way of conclusion, let us begin by recapitulating the analogy between classical statistical mechanics (CSM) and extended probability ensemble quantum theory with one real history (EPE).

There is a common central idea in CSM and EPE: They are both concerned with systems with one real fine-grained history about which we have little information from observation either in practice (CSM) or in principle (EPE). The coarse-grained regularities that are accessible to observation and test cannot

²⁰This was pointed out to us by Dowker, who has worked out specific examples in simple models [31].

²¹Indeed, Dowker has exhibited [31] simple examples involving spin in which, for every fine-grained history, there is some coarse graining in which its probability is zero.

TABLE I. The ensemble in classical statistical mechanics and quantum mechanics.

	Classical statistical mechanics	Quantum mechanics
Real fine-grained history	A particular path in phase space, $z(t)$, obeying an equation of motion	A particular path in configuration space, $q(t)$ between t_0 and t_f
Ensemble	Alternative phase space paths	Alternative configuration space paths between t_0 and t_f
Betting instructions	Using probabilities	Using extended probabilities—the instruction is “don’t bet” if nonstandard probabilities involved
State	Distribution on phase space $\rho(z_0, t_0)$	Wave function $\hat{\Psi}(q_0, t_0)$
Fundamental distribution	$w[z(t)] \equiv \int dz_0 \delta[z(t) - z_t(z_0)] \rho(z_0)$, where $z_t(z_0)$ is the classically evolved z_0	$w[q(t)] \equiv \text{Re}[\hat{\Psi}^*(q_f, t_f) \exp\{iS[q(t)]/\hbar\} \hat{\Psi}(q_0, t_0)]$
Coarse graining	Partitions of the ensemble into classes c_α (coarse-grained histories) one of which contains the real fine-grained history	Partitions of the ensemble into classes c_α (coarse-grained histories) one of which contains the real fine-grained history
Probabilities or extended probabilities for coarse-grained histories	Sum over fine-grained probabilities $p(\alpha) = \int_{c_\alpha} \delta z w[z(t)]$	Sum over fine-grained extended probabilities $p(\alpha) = \int_{c_\alpha} \delta q w[q(t)]$
Sets that are the basis of settleable bets	All coarse-grained sets, in principle	Recorded coarse-grained sets

therefore be predicted from a fine-grained starting point. Rather, both theories use the ensemble method to describe coarse-grained regularities. The real history is conceptually embedded in an ensemble of comparable imagined histories that are assigned measures of ignorance consistent with coarse-grained knowledge.²² Within this ensemble framework there are similarities and differences between CSM and EPE. Table I summarizes these, but we now describe them in a few additional words.

Notion of probability. In CSM the fine-grained history is assumed to be determinable in principle and therefore the basis of a settleable bet on what it is like. Standard probabilities are therefore assigned to members of the ensemble. In EPE the fine-grained history is in principle not completely accessible and therefore not the basis of a settleable bet. The ensemble therefore has to be constructed with extended probabilities.

Assignment. In CSM, for the case of local equilibrium, the probabilities can be constructed by a principle of maximizing entropy, the measure of ignorance, while holding certain quantities fixed. (Energy, momentum, and particle number in suitably small volumes are examples.) In EPE the extended probabilities are given as the fundamental distribution (2.2) based on the state and Hamiltonian. Whether there are deeper principles that underlie this formula is an open question.

Despite these differences the two theories are similar in their general characteristics. In particular, EPE can be regarded as a classical stochastic theory based on extended probabilities. That perspective has a number of advantages for interpreting quantum mechanics that we have discussed earlier: EPE provides a unified perspective on coarse graining. EPE allows the use of ordinary language especially with existential words such as “happen.” EPE provides a simple

interpretation of extended probabilities as a unified measure both of ignorance and knowability in principle. The sum rules of probability theory are satisfied exactly for the extended probabilities of EPE. EPE may provide a starting point for further generalization and modification of quantum mechanics. But perhaps most interestingly, EPE provides an intuitively attractive notion of reality in quantum theory.

If a notion of reality is to be introduced in DHQM it seems only natural that there be one real history in each realm. The question of the connection between the real histories in different realms then arises.²³ Two realms may be compatible in the sense that there is a finer-grained realm of which they are both coarse grainings. In that case the real history in each of the coarser-grained realms is the one that contains the real history of the finer-grained realm. However, two realms may be incompatible—without a finer-grained realm of which they are both coarse grainings. Unadorned DHQM provides no connection between the meanings of reality for incompatible realms. This poses a challenge to the notion of reality in unadorned DHQM.

This challenge is overcome in the EPE version of DHQM. There is one real fine-grained history in an ensemble of comparable histories with extended probabilities. All realms are coarse grainings of this unique fine-grained set described in sum-over-histories variables. The real history in each realm is the one that contains the real fine-grained history. Thus EPE has a single notion of reality that is expressed in all realms.

ACKNOWLEDGMENTS

We thank Fay Dowker for a helpful exchange. J.B.H. thanks the Santa Fe Institute for supporting many visits there. The work of J.B.H. was also supported in part by the National

²²Gibbs considered a real system (box of gas) embedded in an ensemble of imagined systems. In what amounts to the same thing, we are considering an ensemble of real and imagined histories of one system.

²³There is a significant literature discussing reality in the context of decoherent histories quantum mechanics. Some recent discussions can be found in Refs. [32–34] with references to further literature.

Science Foundation under Grant No. PHY07-57035. M.G.-M. and J. B. H. acknowledge the support of the Bryan J. and June B. Zwan Foundation. M.G.-M. was in addition supported by the C.O.U.Q. Foundation, by Insight Venture Partners, and by the KITP in Santa Barbara. The generous help provided by these organizations is gratefully acknowledged by both authors.

APPENDIX: QUESTIONS THAT MIGHT BE FREQUENTLY ASKED

Bell's inequalities. Since EPE can be thought of as a classical stochastic theory, won't it imply that the Bell inequalities are satisfied? As discussed in the introduction, it is well known [13] that to understand classically the experiments that demonstrate a violation of Bell's inequalities requires either nonlocal interactions or negative probabilities or both. EPE can be viewed as a classical stochastic theory *with extended probabilities that can sometimes be negative*. There is thus no conflict with these experiments. Indeed, EPE is a formulation of quantum theory, so it predicts a violation of the Bell inequalities.

Hidden variables. Is this in effect a hidden-variable theory? There are no variables involved beyond the usual quantum fields of sum-over-histories quantum theory—the $\{q(t)\}$. However, their fine-grained values are not completely accessible to experiment or observation and therefore partially hidden.

One real history in each realm. Why are extended probabilities needed? Couldn't one say that in DHQM one history in each realm is real? One can say that. But, as discussed in Sec. VI, using extended probability for fine-grained histories provides a connection between the realities of different realms.

Probabilities and frequencies. I understand probabilities as frequencies, but a negative number cannot be a frequency. We are taking a more general view of probabilities, commonly called a Bayesian one, of probabilities as instructions for betting. Extended probabilities outside the range $[0,1]$ are, in a way, instructions for betting—do not bet, it will not be settleable. The connection of standard probabilities with frequencies of occurrences in an infinite ensemble of identical systems can be derived from this view of probability for realms (see, e.g., [35]).

-
- [1] R. B. Griffiths, *Consistent Quantum Theory* (Cambridge University Press, Cambridge, 2002).
 - [2] R. Omnès, *Interpretation of Quantum Mechanics* (Princeton University Press, Princeton, 1994).
 - [3] M. Gell-Mann, *The Quark and the Jaguar* (W. H. Freeman, New York, 1994).
 - [4] J. B. Hartle, *The Quantum Mechanics of Closed Systems*, Directions in General Relativity, Vol. 1, edited by B.-L. Hu, M. P. Ryan, and C. V. Vishveshwara (Cambridge University Press, Cambridge, 1993).
 - [5] J. B. Hartle, *Spacetime Quantum Mechanics and the Quantum Mechanics of Spacetime*, Gravitation and Quantizations, Proceedings of the 1992 Les Houches Summer School, edited by B. Julia and J. Zinn-Justin, Les Houches Summer School Proceedings, Vol. LVII, (North Holland, Amsterdam, 1995); A précis of these lectures is given in J. B. Hartle, e-print [arXiv:gr-qc/9508023](http://arxiv.org/abs/gr-qc/9508023).
 - [6] J. Willard Gibbs, *Elementary Principles in Statistical Mechanics* (Yale University Press, New Haven, 1902).
 - [7] M. Gell-Mann and S. Lloyd, *Effective Complexity*, in Non-Extensive Entropy—Interdisciplinary Applications, edited by M. Gell-Mann and C. Tsallis (Oxford University Press, New York, 2004).
 - [8] J. B. Hartle, *Phys. Rev. A* **78**, 012108 (2008).
 - [9] M. Gell-Mann and J. B. Hartle, *Phys. Rev. A* **76**, 022104 (2007).
 - [10] J. B. Hartle, *Vistas Astron.* **37**, 569 (1993).
 - [11] M. Gell-Mann, *Physics Today* **42**, 50 (1989).
 - [12] J. B. Hartle, *Generalizing Quantum Mechanics for Quantum Spacetime*, The Quantum Structure of Space and Time, *Proceedings of the 23rd Solvay Conference on Physics*, edited by D. Gross, M. Henneaux, and A. Sevrin (World Scientific, Singapore, 2007).
 - [13] See, e.g., R. P. Feynman, *Negative Probability*, Quantum Implications: Essays in Honor of David Bohm, edited by B. J. Hiley and F. D. Peat (Routledge & Kegan Paul, London, 1987); K. Wódkiewicz, *Phys. Lett. A* **129**, 1 (1988); E. C. G. Sudarshan and T. Rothman, *Int. J. Theor. Phys.* **32**, 1077 (1993); Y. D. Han, W. Y. Hwang, and I. G. Koh, *Phys. Lett. A* **221**, 283 (1996).
 - [14] D. Bohm and B. J. Hiley, *The Undivided Universe* (Routledge, London, 1993).
 - [15] R. P. Feynman, and A. Hibbs, *Quantum Mechanics and Path Integrals* (McGraw-Hill, New York, 1965).
 - [16] B. de Finetti, *Ann. Inst. Henri Poincaré*, **7**, 1 (1937); [English translation *Foresight: Its Logical Laws, its Subjective Sources*, Studies in Subjective Probability, edited by H. E. Kyburg Jr. and H. E. Smokler (John Wiley, New York, 1964)].
 - [17] C. M. Caves, *Probabilities as Betting Odds and the Dutch Book* (unpublished) [<http://info.phys.unm.edu/~caves/reports/reports.html>].
 - [18] J. B. Hartle, *Phys. Rev. A* **70**, 022104 (2004).
 - [19] J. B. Hartle and S. W. Hawking, *Phys. Rev. D* **28**, 2960 (1983).
 - [20] M. Gell-Mann and J. B. Hartle, *Phys. Rev. D* **47**, 3345 (1993).
 - [21] S. Goldstein and D. N. Page, *Phys. Rev. Lett.* **74**, 3715 (1995).
 - [22] C. Caves, *Phys. Rev. D* **33**, 1643 (1986).
 - [23] E. Joos and H. D. Zeh, *Zeit. Phys. B* **59**, 223 (1985).
 - [24] M. Gell-Mann and J. B. Hartle, *Quantum Mechanics in the Light of Quantum Cosmology*, Complexity, Entropy, and the Physics of Information, SFI Studies in the Sciences of Complexity, Vol. VIII, edited by W. Zurek (Addison Wesley, Reading, 1990).
 - [25] M. Gell-Mann and J. B. Hartle, e-print [arXiv:gr-qc/9404013](http://arxiv.org/abs/gr-qc/9404013).
 - [26] M. Gell-Mann and J. B. Hartle, *Time Symmetry and Asymmetry in Quantum Mechanics and Quantum Cosmology*, The Physical Origins of Time Asymmetry, edited by J. Halliwell, J. Pérez-Mercader, and W. Zurek (Cambridge University Press, Cambridge, 1994); e-print [arXiv:gr-qc/9309012](http://arxiv.org/abs/gr-qc/9309012); for some other discussions of time neutrality in quantum theory see, e.g., [1,36].
 - [27] L. Diósi, *Phys. Rev. Lett.* **92**, 170401 (2004).
 - [28] J. B. Hartle, *J. Phys. A: Math. Theor.* **40**, 3101 (2007).

- [29] J. B. Hartle, *Am. J. Phys.* **73**, 101 (2005).
- [30] J. B. Hartle and T. Hertog, e-print [arXiv:1104.1733](#).
- [31] F. Dowker (private communication); see also F. Dowker and Y. Ghazi-Tabatabai, *J. Phys. A* **41**, 105301 (2008).
- [32] *Many Worlds? Everett, Quantum Theory, and Reality*, edited by S. Saunders, J. Barrett, A. Kent, and D. Wallace (Oxford University Press, Oxford, 2010).
- [33] R. Griffiths, e-print [arXiv:1105.3932](#).
- [34] P. C. Hohenberg, e-print [arXiv:1101.2635](#).
- [35] J. B. Hartle, *Am. J. Phys.* **36**, 704 (1968); E. Farhi, J. Goldstone, and S. Gutmann, *Ann. Phys. (NY)* **192**, 368 (1989); Y. Ohkuwa, *Phys. Rev. D* **48**, 1781 (1993).
- [36] Y. Aharonov, P. Bergmann, and J. Lebovitz, *Phys. Rev. B* **134**, 1410 (1964); Y. Aharonov and L. Vaidman, *The Two-State Vector Formalism: An Updated Review*, Time in Quantum Mechanics, edited by J. G. Muga, R. Sala Mayato, and Í. L. Egusquiza, *Lect. Notes Phys.* **734** (Springer, Berlin, 2008); L. S. Schulman, *Time's Arrows and Quantum Measurement* (Cambridge University Press, Cambridge, 1997).